

University of Groningen

## Functionalization of DNA by electrostatic bonding

Chen, Wei

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Chen, W. (2019). *Functionalization of DNA by electrostatic bonding*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Propositions  
Belonging to the dissertation

Functionalization of DNA by Electrostatic Bonding

1. DNA-lipid exchange with amine derived molecules can be achieved stoichiometrically in the organic phase. (Chapter 2)
2. DNA can act as a scaffold for the self-assembly of bound lipids to realize new functions, like a multi-chromophoric light harvesting system. (Chapter 2)
3. Cyclodextrin can be transformed into an ionic liquid with moderate fluidic property at room temperature through a lipid exchange process. (Chapter 3)
4. The approach of DNA-lipid exchange is also applicable to macromolecules like polyethylene glycol (PEG). The degree of exchange is decreasing with the increase of PEG molecular weight. (Chapter 4)
5. Quaternary ammonium-lipids can be exchanged onto DNA by introducing the counterion, acetylacetate, onto these lipids. The acetylacetate can abstract a proton from DNA-ANI complex to facilitate the binding between negatively charged DNA and positively charged quaternary ammonium-lipids. (Chapter 5)
6. DNA bonded with the quaternary ammonium-PEG is more vulnerable to ion displacement than that of primary amine-PEG. (Chapter 5)
7. The development of proper characterization methods to fully understand the conformation of DNA-lipids is challenging.
8. Brave is more crucial than clever in scientific research.